

## HPLC Analysis of Oligonucleotides Using YMC's Non-porous AEX Columns

Oligonucleotides are important in genetic testing, research and forensics. For quite some time now, oligonucleotide-based approaches have been developed for different pharmaceutical applications.

Therefore, corresponding analytical methods have to be available. In addition to RP based methods, IEX methods can provide an alternative approach. The

strong anion exchanger BioPro IEX QF has a quaternary amine residue as the functional group. The non-porous BioPro IEX QF offers high efficiency, exceptionally high resolution at low operating pressures.

**A method for the determination of oligonucleotides using BioPro IEX QF has been developed by YMC. 3 application examples are presented here.**

Table 1: Chromatographic conditions

Column:	BioPro IEX QF (5 $\mu$ m) 100 x 4.6 mm ID	
Part No.:	QF00S05-1046WP	
Eluent:	A) 10 mM NaOH B) 10 mM NaOH containing 1.0 M NaClO <sub>4</sub>	
Gradient:	Time [min]	Eluent B [%]
	0	25
	15	55
	20	100
Flow rate:	1.0 mL/min	
Temperature:	25 °C	
Detection:	UV at 260 nm	
Injection:	4 $\mu$ L (5 nmol/mL)	




Figure 1: Separation of synthetic oligonucleotides (Single-strand DNA)





Figure 2: Separation of synthetic oligonucleotides (Single-strand RNA)




Figure 3: Separation of synthetic oligonucleotides (Single-strand RNA)

